



EJVES EXTRA ABSTRACT[☆]

Venous Aneurysms – An Uncommon but Potentially Life-Threatening Disease: A Case Report

D. Papadimitriou, M. Tachtsi, G. Pitoulas, S. Atmatzidis, P. Fycatas, D. Christopoulos
Aristotle University of Thessaloniki, 2nd Surgical Department, Division of Vascular Surgery, General Hospital "G.Gennimatas," Thessaloniki, Greece

Venous aneurysms are rare, but they can have devastating consequences, including pulmonary embolism (PE) and death. We present a case of a male patient with venous aneurysms of both his legs treated surgically.

Patient and methods: A 24-year-old patient appeared with varicosities of both his legs. Magnetic resonance venography (MRV) revealed partial thrombosis of the vena cava and venous aneurysms of the left popliteal and calf vein and also of the right small saphenous vein (SSV). The patient was treated with tangential excision and lateral venoraphy of the popliteal venous aneurysm (PVA), resection and ligation of the aneurysms of the left SSV with an excellent result.

Conclusion: Our results are in agreement with the literature. Surgical resection is recommended whenever possible, because there is a significant risk of pulmonary embolism and death if left untreated.

doi:10.1016/j.ejvs.2009.04.010

DOI of original article:10.1016/j.ejvsextra.2009.04.003

Available online 27 August 2009

An Unusual Case of Post-stent Carotid Artery Stenosis with Successful Operative Repair: A Case Report

H. Mistry, A. Currie, C. Lioupis, M. Tyrrell, D. Valenti
Department of Vascular Surgery, King's College London, Denmark Hill, London SE5 9RS, UK

We report a symptomatic internal carotid artery re-stenosis distal to a carotid stent in a 69-year-old man, which resulted in a stroke 6 months post-stenting. It was treated by a successful carotid endarterectomy and explantation of the stent.

doi:10.1016/j.ejvs.2009.07.018

DOI of original article:10.1016/j.ejvsextra.2009.07.005

Available online 29 August 2009

Endovascular Treatment for Left-side Cervical Aortic Arch with Huge Aneurysm

W. Hai-tao^a, M. Jun^a, J. Wei-liang^b, W. Qing-hua^a
^a *Department of Vascular Surgery, Capital Medical University Affiliated Anzhen Hospital, Beijing 100029, China*
^b *Department of Vascular Surgery, Harbin Medical University Affiliated 2nd Hospital, Harbin, Heilongjiang 150086, China*

Cervical aortic arch (CAA) is an unusual malformation of the aortic arch. CAA with aneurysm formation is very rare. A 67-year-old female patient was admitted with a huge mass on left-side neck and 2 days of dyspnea. Investigations revealed a huge aneurysm on a left-side CAA. The aneurysm was successfully repaired using thoracic aortic stent-graft. The patient's dyspnea was resolved immediately.

doi:10.1016/j.ejvs.2009.07.022

DOI of original article:10.1016/j.ejvsextra.2009.07.004

Available online 26 August 2009

[☆] Full articles available online at www.ejvsextra.com